

***Salmonella enterica* subsp. *enterica*, Strain LS1059**

Catalog No. NR-4347

For research use only. Not for human use.

Contributor:

National Institute of Allergy and Infectious Diseases, National Institutes of Health

Manufacturer:

BEI Resources

Product Description:

Bacteria Classification: *Enterobacteriaceae*, *Salmonella*

Species: *Salmonella enterica*

Subspecies: *Salmonella enterica* subsp. *enterica*^{1,2}

Serogroup: B

Serovar: Typhimurium

Strain: LS1059

Original Source: *Salmonella enterica* (*S. enterica*) subsp. *enterica*, strain LS1059 is a derivative of strain 14028.³ Strain 14028 (also referred to as CDC 6516-60) was isolated from chickens and deposited to the ATCC® in 1960.

Comments: *S. enterica* subsp. *enterica*, strain LS1059 expresses STM3119, tagged with 3 X FLAG epitopes.³ STM3119 is proposed to function as a monoamine oxidase (MaoA) that converts aminoacetone, which is the degradation product of L-threonine, to methylglyoxal.³ Additional information is available at the [Resource Center for Biodefense Proteomics Research \(BPRC\)](#).

S. enterica are Gram-negative, rod-shaped, flagellated bacteria. The species is divided into six subspecies (I, II, IIIa, IIIb, IV, VI) where only subspecies I, subsp. *enterica*, is considered of clinical relevance. Salmonellosis (non-typhoidal), due to the greater than 1500 serovars of *S. enterica* subsp. *enterica*, is one of the most common food-borne diseases with an estimated 2 million cases that occur in the United States every year.⁴ Pathogenicity results from a variety of virulence factors found in plasmids, prophages, and five pathogenicity islands which allow these organisms to colonize and infect host organisms.^{5,6}

S. enterica subsp. *enterica* serovar Typhimurium (formerly *Salmonella typhimurium*) is a major cause of gastroenteritis. These bacteria are host generalists that occur in humans and many other mammals. Septic shock resulting in part from lipopolysaccharide (LPS) is a primary complication associated with serovar Typhimurium infection.⁷ Due to its similarity to the clinical and pathological effects in humans, calves are currently used as an animal model for human enterocolitis caused by this serotype.⁸ Additionally, this serovar causes typhoid-like disease in mice and is used as a mouse model of human typhoid fever.⁹

The complete genome sequence of several strains of *S. enterica* subsp. *enterica* serovar Typhimurium are in progress [strain DT104 (Definitive Type 104; a multidrug resistant strain), strain SL1344 (a genetically marked subline of a calf-virulent isolate), and strain TR7095 (a wild-type strain)] and strain LT2 has been completed (GenBank: AE006468).⁹

Material Provided:

Each vial contains approximately 0.5 mL of bacterial culture in 0.5X Tryptic Soy Broth supplemented with 10% glycerol.

Note: If homogeneity is required for your intended use, please purify prior to initiating work.

Packaging/Storage:

NR-4347 was packaged aseptically, in screw-capped plastic cryovials. The product is provided frozen and should be stored at -60°C or colder immediately upon arrival. For long-term storage, the vapor phase of a liquid nitrogen freezer is recommended. Freeze-thaw cycles should be avoided.

Growth Conditions:

Media:

Tryptic Soy Broth or equivalent

Tryptic Soy Agar or equivalent

Incubation:

Temperature: 37°C

Atmosphere: Aerobic

Propagation:

1. Keep vial frozen until ready for use, then thaw.
2. Transfer the entire thawed aliquot into a single tube of broth.
3. Use several drops of the suspension to inoculate an agar slant and/or plate.
4. Incubate the tubes and plate at 37°C for 24 hours.

Citation:

Acknowledgment for publications should read "The following reagent was obtained through BEI Resources, NIAID, NIH: *Salmonella enterica* subsp. *enterica*, Strain LS1059, NR-4347."

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. Biosafety in Microbiological and Biomedical Laboratories, 5th ed. Washington, DC: U.S. Government Printing Office, 2007; see www.cdc.gov/od/ohs/biosfty/bmbl5/bmbl5toc.htm.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at www.beiresources.org.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government make any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, non-commercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

References:

- Judicial Commission of the International Committee on Systematics of Prokaryotes. "The Type Species of the Genus *Salmonella* Lignierres 1900 Is *Salmonella enterica* (ex Kauffmann and Edwards 1952) Le Minor and Popoff 1987, with the Type Strain LT2^T, and Conservation of the Epithet *enterica* in *Salmonella enterica* over All Earlier Epithets that May Be Applied to This Species. Opinion 80." Int. J. Syst. Evol. Microbiol. 55 (2005): 519-520. PubMed: 15653929.
- Tindall, B. J., et al. "Nomenclature and Taxonomy of the Genus *Salmonella*." Int. J. Syst. Evol. Microbiol. 55 (2005): 521-524. PubMed: 15653930.
- Shi, L., et al. "Proteomic Analysis of the *Salmonella enterica* Serovar Typhimurium Isolated from RAW 264.7 Macrophages: Identification of a Novel Protein that Contributes to the Replication of Serovar Typhimurium Inside Macrophages." J. Biol. Chem. 281 (2006): 29131-29140. PubMed: 16893888.
- Altekruse, S. F., M. L. Cohen and D. L. Swerdlow. "Emerging Foodborne Diseases." Emerg. Infect. Dis. 3 (1997): 285-293. PubMed: 9284372.
- Lavigne, J. P. and A. B. Blanc-Potard. "Molecular Evolution of *Salmonella enterica* Serovar Typhimurium and Pathogenic *Escherichia coli*: From Pathogenesis to Therapeutics." Infect. Genet. Evol. 8 (2008): 217-226. PubMed: 18226587.
- Parsons, D. A. and F. Heffron. "*sciS*, an *icmF* Homolog in *Salmonella enterica* Serovar Typhimurium, Limits Intracellular Replication and Decreases Virulence." Infect. Immun. 73 (2005): 4338-4345. PubMed: 15972528.
- Sha, J., et al. "The Two Murein Lipoproteins of *Salmonella enterica* Serovar Typhimurium Contribute to the Virulence of the Organism." Infect. Immun. 72 (2004): 3987-4003. PubMed: 15213144.
- Zhang, S., et al. "The *Salmonella enterica* Serotype Typhimurium Effector Proteins SipA, SopA, SopB, SopD, and SopE2 Act in Concert to Induce Diarrhea in Calves." Infect. Immun. 70 (2002): 3843-3855. PubMed: 12065528.
- McClelland, M., et al. "Complete Genome Sequence of *Salmonella enterica* Serovar Typhimurium LT2." Nature 413 (2001): 852-856. PubMed: 11677609. GenBank: AE006468.

ATCC® is a trademark of the American Type Culture Collection.

