



Product Information Sheet for HRP-20122

Simian Immunodeficiency Virus, SIVsmE660-807-24w Env

Catalog No. HRP-20122

This reagent is the tangible property of the U.S. Government.

Lot No. 70052311

For research use only. Not for use in humans.

Contributor and Manufacturer:

Vanessa M. Hirsch, D.V.M., D.Sc., Laboratory of Molecular Biology, National Institute of Allergy and Infectious Diseases (NIAID), National Institutes of Health (NIH), Bethesda, Maryland, USA

Product Description:

VIRUS CLASSIFICATION: *Retroviridae, Lentivirus*

SPECIES: Simian immunodeficiency virus

STRAIN/ISOLATE: SIVsmE660-807-24w Env

ORIGINAL SOURCE: Simian immunodeficiency virus (SIV), SIVsmE660-807-24w Env is a derivative of SIVsmE660-FL14 (HRP-20120), encoding envelope glycoprotein ([JQ864168](#)) cloned from neutralizing antibody escape variant obtained at 24 weeks post-infection from a SIVsmE660-FL14-inoculated rhesus macaque (*Macaca mulatta*).^{1,2,3}

COMMENTS: SIVsmE660-807-24w Env exhibits Tier 3 neutralization antibody resistance.² The complete genome of the SIVsmE660-FL14 clone has been sequenced (GenBank: [JQ864087.1](#)).¹

Material Provided:

Each vial contains approximately 0.5 mL of supernatant from rhesus macaque peripheral blood leukocytes (PBL) infected with SIVsmE660-807-24w Env. The virus supernatants were prepared by centrifugation followed by filtration through a 0.45 µm filter. The TCID₅₀ titer in TZM-bl cells was 51,200 infectious units (IU) per mL. HRP-20122 has not been tested for mycoplasma contamination.¹

Note: If homogeneity is required for your intended use, please purify prior to initiating work.

Packaging/Storage:

HRP-20122 was packaged aseptically in plastic cryovials. The product is provided frozen and should be stored at -100°C or colder immediately upon arrival. For long-term storage, the vapor phase of a liquid nitrogen freezer is recommended. Freeze-thaw cycles should be avoided.

Growth Conditions:

HOST: Rhesus macaque peripheral blood mononuclear cells (PBMC)

GROWTH MEDIUM: RPMI 1640 medium supplemented with 10% heat-inactivated fetal bovine serum

INFECTION: Cells should be 70% to 90% confluent

INCUBATION: 10 to 14 days at 37°C and 5% CO₂

Citation:

Acknowledgment for publications should read "The following reagent was obtained through the NIH HIV Reagent Program, NIAID, NIH: Simian Immunodeficiency Virus, SIVsmE660-807-24w Env, HRP-20122, contributed by Dr. Vanessa M. Hirsch."

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. Biosafety in Microbiological and Biomedical Laboratories. 6th ed. Washington, DC: U.S. Government Printing Office, 2020; see www.cdc.gov/biosafety/publications/bmbl5/index.htm.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the NIH HIV Reagent Program Material Transfer Agreement (MTA). The MTA is available on our Web site at www.hivreagentprogram.org.

While the NIH HIV Reagent Program uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government makes any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use, and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure the authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to the NIH HIV Reagent Program are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, non-commercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a

NIH HIV Reagent Program

www.hivreagentprogram.org

E-mail: contact@HIVReagentProgram.org

Tel: 888-487-0727 | Fax: 703-365-2898

© 2022 American Type Culture Collection (ATCC)

All rights reserved.

HRP-20122_02SEP2022

Page 1 of 2



Product Information Sheet for HRP-20122

license is required. U.S. Government contractors may need a license before first commercial sale.

References:

1. Hirsch, V. M., Personal Communication.
2. Wu, F., et al. "SIV Infection Duration Largely Determines Broadening of Neutralizing Antibody Response in Macaques." *J. Clin. Invest.* 130 (2020): 5413-5424. PubMed: 32663192.
3. Wu, F., et al. "Sequential Evolution and Escape from Neutralization of Simian Immunodeficiency Virus SIVsmE660 Clones in Rhesus Macaques." *J. Virol.* 86 (2012): 8835-8847. PubMed: 22696650.

ATCC® is a trademark of the American Type Culture Collection.



NIH HIV Reagent Program

www.hivreagentprogram.org

E-mail: contact@HIVReagentProgram.org

Tel: 888-487-0727 | Fax: 703-365-2898

© 2022 American Type Culture Collection (ATCC)

All rights reserved.

HRP-20122_02SEP2022

Page 2 of 2