

**Dengue Virus Type 2, K0049**

**Catalog No. NR-12215**

**Product Description:** Cell lysate and supernatant from *Aedes albopictus* clone C6/36 cells<sup>1</sup> infected with dengue virus type 2 (DEN-2), K0049.

**Lot<sup>2</sup>: 58521715**

**Manufacturing Date: 17JUL2009**

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity in C6/36 Cells <sup>1</sup>	Report results	Sloughing of cells
Identification by Indirect Fluorescent Antibody (IFA) Assay <sup>3</sup>	Fluorescence observed	Fluorescence observed
Sequencing of DEN-2 Specific Sequence (963 nucleotides)	Consistent with DEN-2	Consistent with DEN-2
Titer by TCID <sub>50</sub> Assay in C6/36 Cells with IFA Readout <sup>1,4,5</sup>	Report results	2.81 x 10 <sup>7</sup> TCID <sub>50</sub> /mL
Functional Activity by RT-PCR Assay Using DEN-2 Specific Primers	~ 1200 bp amplicon	~ 1200 bp amplicon
<b>Sterility (21-day incubation)</b> Harpo's HTYE broth <sup>6</sup> , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Brucella agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO <sub>2</sub>	No growth No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth No growth
<b>Mycoplasma Contamination</b> Agar and broth culture (14-day incubation at 37°C) DNA Detection by PCR of Test Article nucleic acid	None detected None detected	None detected None detected

<sup>1</sup>*Aedes albopictus* clone C6/36 cells (ATCC® CRL-1660™)

<sup>2</sup>DEN-2, K0049 was deposited by Dr. Rebeca Rico-Hesse of the Department of Virology and Immunology, Southwest Foundation for Biomedical Research, San Antonio, Texas. NR-12215 was grown from deposited virus seed in Minimum Essential Medium containing Earle's salts and non-essential amino acids (Invitrogen™ 10370) supplemented with 2% fetal bovine serum (ATCC® 30-2020), 2 mM L-glutamine (Invitrogen™ 25030-081), and 1 mM sodium pyruvate (Invitrogen™ 11360-070) for 7 days at 28°C with 5% CO<sub>2</sub>

<sup>3</sup>Using monoclonal antibody specific to DEN-2 (Chemicon MAB8702)

<sup>4</sup>The Tissue Culture Infectious Dose 50% (TCID<sub>50</sub>) endpoint is the 50% infectious endpoint in cell culture. The TCID<sub>50</sub> is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD<sub>50</sub>) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID<sub>50</sub> provides a measure of the titer (or infectivity) of a virus preparation.

<sup>5</sup>7 days at 28°C with 5% CO<sub>2</sub>

<sup>6</sup>Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

**Date:** 29 MAR 2010

**Signature:** Signature on File

**Title:** Technical Manager, BEI Authentication or designee

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

