

Certificate of Analysis for HM-654

Bacteriophage Phi TK1718sp/PAK, Infectious for Pseudomonas aeruginosa

Catalog No. HM-654

Product Description: Bacteriophage phi_TK1718sp/PAK was isolated from TK1718 bacterium cultured from leg, foot or decubitus ulcers of a human patient. Bacteriophage phi TK1718sp/PAK (HMP ID 9764) is a reference genome for The Human Microbiome Project (HMP).

Lot^{1,2}: 59978206 Manufacturing Date: 13JUL2011

TEST	SPECIFICATIONS	RESULTS
Plaque Morphology ³	Report results	Circular and clear
Titer ³ Plaque-forming units (pfu) with Pseudomonas aeruginosa	>10 ⁶ pfu per mL	8 x 10 ¹⁰ pfu per mL
Sterility	0.22 µm filtered	0.22 µm filtered
Bacterial Inactivation 10% of total yield plated on Tryptic Soy Agar with 5% defibrinated sheep blood ⁴	No viable bacteria detected	No viable bacteria detected

¹Quality control of HMP material is only performed to demonstrate that the product distributed by BEI Resources is identical to the deposited material. It should not be considered a complete characterization of the deposited material.

Date: 09 APR 2013 Signature:

> Technical Manager, BEI Authentication or designee Title:

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

E-mail: contact@beiresources.org **BEI Resources** www.beiresources.org

Tel: 800-359-7370 Fax: 703-365-2898

²The deposited material was added to a Luria-Burtani (LB) soft agar overlay (0.5%) supplemented with 10 mM MgSO₄ containing the phage host, Pseudomonas aeruginosa, strain PAK. The soft agar overlay was added to an LB agar Kolle supplemented with 10 mM MgSO₄ and incubated 24 hours at 30°C in an aerobic atmosphere. After two additional passages under the above conditions, the bacteriophage were harvested to produce this lot.

³Lysis was observed on the soft agar overlay after 24 hours at 30°C in an aerobic atmosphere.

⁴7 days at 37°C in an aerobic atmosphere