

Venezuelan Equine Encephalitis Virus, MX03-H1

Catalog No. NR-21709

Product Description: Cell lysate and supernatant from *Cercopithecus aethiops* kidney epithelial cells (Vero)¹ infected with Venezuelan equine encephalitis virus (VEEV), MX03-H1

Lot²: 60858747

Manufacturing Date: 11JUN2012

| TEST | SPECIFICATIONS | RESULTS |
|--|--|--|
| Identification by Infectivity Using Vero Cells ¹ | Report results | Rounding and detachment |
| Sequencing of Species-Specific Region (699 nucleotides) | Consistent with VEEV, MX03-H1 | 100% identity with VEEV, MX03-H1 (GenBank: KC344482) |
| Titer by TCID ₅₀ Assay ^{3,4} in Vero Cells ¹ | Report results | 8.9 × 10 ⁸ TCID ₅₀ per mL |
| Functional Activity by RT-PCR Assay | ~ 750 bp amplicon | ~ 750 bp amplicon |
| Sterility (21-day incubation) Harpo's HTYE broth ⁵ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂ | No growth No growth No growth No growth No growth No growth No growth No growth | No growth No growth No growth No growth No growth No growth No growth No growth |
| Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) | None detected | None detected |

¹Vero cells: ATCC® CCL-81™

²Grown in Eagle's Minimum Essential Medium containing Earle's Balanced Salt Solution, non-essential amino acids, 2 mM L-glutamine, 1 mM sodium pyruvate and 1.5 g/L of sodium bicarbonate (ATCC® 30-2003) supplemented with 2% fetal bovine serum (ATCC® 30-2020) for 3 days at 37°C with 5% CO₂.

³The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the titer (or infectivity) of a virus preparation.

⁴6 days at 37°C and 5% CO₂

⁵Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

Date: 24 MAR 2014

Signature: 

Title: Technical Manager, BEI Authentication or designee

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

