

***Trypanosoma brucei* subsp. *rhodesiense*, Strain KETRI 2482**

Catalog No. NR-46434

Product Description: *Trypanosoma brucei* (*T. brucei*) subsp. *rhodesiense*, strain KETRI 2482 was isolated in 1969 from the blood of a patient before chemotherapy in Lumino, Uganda. *T. brucei* subsp. *rhodesiense*, strain KETRI 2482 was obtained by Professor C. J. Bacchi from the Kenya Trypanosomiasis Research Institute (KETRI) strain bank at Mugaga, Kenya.

Lot¹: 63717831

Manufacturing Date: 04SEP2015

TEST	SPECIFICATIONS	RESULTS
Genotyping Sequencing of internal transcribed spacer (ITS) 1, 5.8S ribosomal RNA gene, ITS 2 (~ 130 base pairs)	Consistent with <i>T. brucei</i>	Consistent with <i>T. brucei</i> ²
Functional Activity by PCR Amplification ITS 1, 5.8S ribosomal RNA gene, ITS 2 ³	~ 1300 base pair amplicon	~ 1300 base pair amplicon
Level of Parasitemia (pre-freeze)⁴	≥ 1 × 10 ⁶ parasites per mL	3 × 10 ⁸ parasites per mL
Viability (post-freeze)⁵	Growth in inoculated mouse	Growth in inoculated mouse

¹NR-46434 was produced by inoculation of the deposited material into a BALB/c mouse. Infection was allowed to progress for 2 days until the first peak of parasitemia was reached. Infected blood was collected by orbital bleeding and used to inoculate six BALB/c mice. Infection was allowed to progress for 2 days until the first peak of parasitemia was reached and infected blood was collected by orbital bleeding.

²Also consistent with *T. evansi* and/or *T. equiperdum*, which are putative subspecies of *T. brucei* (Lun, Z.-R., et al. "Trypanosoma brucei: Two Steps to Spread Out from Africa." *Trends Parasitol.* 26 (2010): 424-427. PubMed: 20561822.)

³PCR was performed as described in Agbo, E. C., et al. "Measure of Molecular Diversity within the *Trypanosoma brucei* Subspecies *Trypanosoma brucei brucei* and *Trypanosoma brucei gambiense* as Revealed by Genotypic Characterization." *Exp. Parasitol.* 99 (2001): 123-131. PubMed: 11846522.

⁴Parasitemia was determined after 2 days of infection by microscopic counts using a haemocytometer and 0.85% ammonium chloride as diluent.

⁵Viability of trypanosomes was confirmed by examination of a BALB/c mouse for parasitemia at daily intervals for 4 days post-infection.

Date: 30 NOV 2015

Signature:

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

