

West Nile Virus, BA 80900-4 (44)

Catalog No. NR-49915

Product Description: Cell lysate and supernatant from *Cercopithecus aethiops* kidney epithelial cells¹ infected with West Nile virus (WNV), BA 80900-4 (44)

Passage History: V1/V3 (Prior to deposit at BEI Resources/BEI Resources); V# = Number of passages in Vero cells²

Lot³: 63902492

Manufacturing Date: 19APR2016

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity in Vero cells	Cell rounding and detachment	Cell rounding and detachment
Sequencing of Species-Specific Region (843 nucleotides)	Consistent with WNV	Consistent with WNV ⁴
Titer by TCID ₅₀ Assay ^{5,6} in Vero cells ¹	Report results	8.9 × 10 ⁸ TCID ₅₀ per mL
Amplification of WNV Sequence by RT-PCR	~ 950 bp amplicon	~ 950 bp amplicon
Sterility (21-day incubation) Harpo's HTYE broth ⁷ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

¹Vero cells: ATCC® CCL-81™

²The second virus passage at BEI Resources was performed by lipofectamine transfection of extracted viral nucleic acid in order to remove contaminating mycoplasma.

³Grown in Eagle's Minimum Essential Medium containing Earle's Balanced Salt Solution, non-essential amino acids, 2 mM L-glutamine, 1 mM sodium pyruvate and 1.5 g/L of sodium bicarbonate (ATCC® 30-2003) supplemented with 2% fetal bovine serum (ATCC® 30-2020) for 4 days at 37°C with 5% CO₂

⁴Sequence information for WNV, BA 80900-4 (44) is not available in the NCBI database; nucleotide sequence obtained for NR-49915, Lot No. 63902492 is highly similar to numerous WNV strains.

⁵The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the titer (or infectivity) of a virus preparation.

⁶4 days at 37°C and 5% CO₂

⁷Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

Date: 16 NOV 2016

Signature: 

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

