

***Plasmodium falciparum*, Strain FCR-3/Gambia Clone I, Knobby**

Catalog No. MRA-737

Product Description: *Plasmodium falciparum* (*P. falciparum*), strain FCR-3/Gambia Clone I, Knobby is a clone derived from isolate FCR-3/FMG (Gambia) after four years of continuous culture by W. Trager by microscopic selection. MRA-737 was derived from ATCC® 50041™, which was deposited at ATCC® by W. Trager.

Lot¹: 60031006

Manufacturing Date: 09JUN2011

TEST	SPECIFICATIONS	RESULTS
Identification by Giemsa Stain Microscopy^{2,3}	Blood-stage parasites present	Blood-stage parasites present
Level of Parasitemia Pre-freeze ^{4,5} Ring-stage parasitemia Post-freeze ^{2,6} Total parasitemia	Report results ≥ 1%	5-6% 7-8%
Viability^{2,7}	Growth in infected red blood cells	Growth in infected red blood cells
Mycoplasma Contamination² DNA Detection by PCR	None detected	None detected

¹MRA-737 was produced by cultivation of BEI Resources MRA-737 lot 4599776 in fresh human erythrocytes suspended in RPMI 1640 medium, adjusted to contain 10% (v/v) heat-inactivated human serum (pooled Type A), 25 mM HEPES, 2 mM L-glutamine, 4 g/L D-glucose, 0.005 µg/mL hypoxanthine and 2.5 µg/mL gentamicin. The culture was incubated at 37°C in sealed flasks outgassed with blood-gas atmosphere (90% N₂, 5% CO₂, 5% O₂) and monitored for parasitemia daily for 8 days. Every 1 to 3 days, uninfected, leukocyte filtered, Type O erythrocytes in complete culture medium were added dropwise to the culture as needed and monitored for hematocrit.

²Testing completed on vial post-freeze material.

³Blood-stage malaria parasites (rings, trophozoites, schizonts +/- gametocytes) were examined by microscopic Giemsa-stained blood smears of an *in vitro* human blood culture over 3 days.

⁴Testing completed on bulk material prior to vialing and freezing.

⁵Parasitemia was determined after 8 days post infection by microscopic counts of Giemsa-stained blood smears.

⁶Parasitemia was determined after 3 days post infection by microscopic counts of Giemsa-stained blood smears.

⁷Viability was confirmed by examination of infected erythrocytes for parasitemia at 3 days post infection.

/Heather Couch/

Heather Couch

20 DEC 2018

Program Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

